



Inducción Floral	Brote 5 - 10 cm	Flor 30-50%	Flor 70-80%	Crecimiento de fruto 20-25mm	Quiebre de color interno	Pre-Cosecha	Post-Cosecha
Dosis	Dosis	Dosis	Dosis	Dosis	Dosis	Dosis	Dosis
1,5 Kg/100L de agua	2 - 3 Kg/ha	0,5 - L/ha	1 L/ha	1 L/ha	5 L/ha	400cc/100L	200-300 cc /100L de agua
<b>Harvest More®</b> 30-10-10	<b>KEY FEEDS® MICROMIX®</b>	<b>STIMULATE</b> FRUIT SIZER	<b>STIMULATE</b> FRUIT SIZER	<b>STIMULATE</b> FRUIT SIZER	<b>Stoller Color</b>	<b>PHYTOGARD®</b>	<b>NATUR'L ÓLEO®</b>
	+ 0,5 - 1 L/ha	+ 3 L/ha	+ 10 ppm	+ 10 ppm	+ 5 Kg/ha		
	<b>ReZist®</b>	<b>SETT®</b>	<b>N-LARGE®</b>	<b>N-LARGE®</b>	<b>Harvest More®</b> 20-20-20		
	+ 4-6 L/ha		+ 3 L/ha	+ 5 K/ha	*Repetir aplicación a los 10-15 días.		
	<b>Keylate</b> zinc		<b>SETT®</b>	<b>Harvest More®</b> 5-5-45			
	+ 2 - 4 L/ha		+ 5 K/ha				
	<b>StollerMagnesio®</b>		<b>Harvest More®</b> 5-5-45				
	* Repetir aplicación en cada brotación de la temporada						

**RootFEED®**  
Concentrate

10 -15 L /ha cada 12-15 días

